

Amendments to Specification

Please replace paragraph 785 on page 196 with the following paragraph:

[785] The term "heteroaryl" (alone or in combination with another term(s)) means an aromatic heterocyclyl containing from 5 to 14 ring atoms. A heteroaryl may be a single ring or 2 or 3 fused rings. Examples of heteroaryl substituents include 6-membered ring substituents such as pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, and 1,3,5-, 1,2,4-, and 1,2,3-triazinyl; 5-membered ring substituents such as imidazolyl, furanyl, thiophenyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, 1,2,3-, 1,2,4-, 1,2,5-, or 1,3,4-oxadiazolyl and isothiazolyl; 6/5-membered fused ring substituents such as benzothiofuranyl, isobenzothiofuranyl, benzisoxazolyl, benzoxazolyl, purinyl, and anthranilyl; and 6/6-membered fused rings such as quinolinyl, isoquinolinyl, cinnolinyl, and quinazolinyl.

Please replace paragraph 1064 on page 291 with the following paragraph:

[1064] The study of angiogenesis depends on a reliable and reproducible model for the stimulation and inhibition of a neovascular response. The corneal micropocket assay provides such a model of angiogenesis in the cornea of a mouse. *See*, Kenyon, B.M., et al., "A Model of Angiogenesis in the Mouse Cornea", *Investigative Ophthalmology & Visual Science*, Vol. 37, No. 8, pp. 1625-1632 (July 1996).